

## Youth Online: High School YRBS

### Arkansas 2009 and United States 2009 Results

High School Youth Risk Behavior Survey						
Question	Arkansas 2009	United States 2009	p-value	Arkansas 2009 More Likely Than United States 2009	United States 2009 More Likely Than Arkansas 2009	No Difference
<b>Unintentional Injuries and Violence</b>						
<b>Rarely or never wore a bicycle helmet</b> (among students who had ridden a bicycle during the 12 months before the survey)	90.1 (86.5–92.8) 1,021 <sup>†</sup>	84.7 (81.2–87.6) 10,486	0.02	●		
<b>Rarely or never wore a seat belt</b> (when riding in a car driven by someone else)	18.7 (14.4–24.0) 1,678	9.7 (8.2–11.4) 16,220	0.00	●		
<b>Rode with a driver who had been drinking alcohol one or more times</b> (in a car or other vehicle during the 30 days before the survey)	28.3 (24.5–32.4) 1,689	28.3 (26.7–29.9) 16,347	0.99			●
<b>Drove when drinking alcohol one or more times</b> (in a car or other vehicle during the 30 days before the survey)	12.5 (10.0–15.4) 1,642	9.7 (8.7–10.8) 16,121	0.05			●
<b>Carried a weapon on school property on at least 1 day</b> (for example, a gun, knife, or club during the 30 days before the survey)	8.4 (6.6–10.8) 1,656	5.6 (5.0–6.3) 16,256	0.01	●		
<b>Did not go to school because they felt unsafe at school or on their way to or from school on at least 1 day</b> (during the 30 days before the survey)	10.4 (7.6–14.0) 1,682	5.0 (4.3–5.7) 16,371	0.00	●		
<b>Threatened or injured with a weapon on school property one or more times</b> (for example, a gun, knife, or club during the 12 months before the survey)	11.9 (9.4–15.1) 1,689	7.7 (6.9–8.5) 16,367	0.00	●		
<b>In a physical fight on school property one or more times</b> (during the 12 months before the survey)	14.8 (12.3–17.7) 1,633	11.1 (10.0–12.2) 16,089	0.01	●		
<b>Bullied on school property</b> (during the 12 months before the survey)	—	19.9 (18.8–21.1) 15,633	~			
<b>Felt sad or hopeless</b> (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	28.1 (24.5–32.0) 1,661	26.1 (24.8–27.5) 16,232	0.31			●
<b>Seriously considered attempting suicide</b> (during the 12 months before the survey)	18.0 (14.7–21.8) 1,658	13.8 (13.1–14.6) 16,220	0.02	●		

<b>Made a plan about how they would attempt suicide</b> (during the 12 months before the survey)	14.1 (11.5–17.2) 1,663	10.9 (10.0–11.8) 16,213	0.03	●		
<b>Attempted suicide one or more times</b> (during the 12 months before the survey)	12.0 (9.5–15.0) 1,392	6.3 (5.7–7.0) 14,609	0.00	●		
<b>Suicide attempt resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse</b> (during the 12 months before the survey)	4.8 (3.4–6.8) 1,374	1.9 (1.6–2.3) 14,514	0.00	●		
<b>Carried a weapon on at least 1 day</b> (for example, a gun, knife, or club during the 30 days before the survey)	22.9 (19.4–26.9) 1,603	17.5 (16.1–19.0) 16,110	0.01	●		
<b>Carried a gun on at least 1 day</b> (during the 30 days before the survey)	10.7 (8.4–13.5) 1,645	5.9 (5.1–6.9) 15,664	0.00	●		
<b>In a physical fight one or more times</b> (during the 12 months before the survey)	34.7 (30.5–39.1) 1,606	31.5 (30.1–32.9) 16,130	0.15			●
<b>Injured in a physical fight one or more times</b> (injuries had to be treated by a doctor or nurse, during the 12 months before the survey)	6.9 (5.2–9.1) 1,648	3.8 (3.3–4.3) 15,749	0.00	●		
<b>Hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend</b> (during the 12 months before the survey)	16.7 (12.9–21.2) 1,650	9.8 (8.9–10.8) 16,217	0.00	●		
<b>Ever physically forced to have sexual intercourse</b> (when they did not want to)	13.1 (11.0–15.5) 1,671	7.4 (6.7–8.3) 15,735	0.00	●		
<b>Tobacco Use</b>						
<b>Ever tried cigarette smoking</b> (even one or two puffs)	54.9 (50.8–59.0) 1,567	46.3 (43.7–48.9) 15,888	0.00	●		
<b>Smoked a whole cigarette for the first time before age 13 years</b>	17.7 (14.7–21.1) 1,550	10.7 (9.6–11.9) 15,739	0.00	●		
<b>Smoked cigarettes on at least 1 day</b> (during the 30 days before the survey)	20.3 (16.7–24.5) 1,551	19.5 (17.9–21.2) 15,762	0.68			●
<b>Smoked cigarettes on 20 or more days</b> (during the 30 days before the survey)	9.7 (7.0–13.2) 1,551	7.3 (6.4–8.3) 15,762	0.13			●
<b>Smoked more than 10 cigarettes per day</b> (among students who currently smoked cigarettes, on the days they smoked during the 30 days before the survey)	10.9 (7.5–15.6) 296	7.8 (6.6–9.0) 2,940	0.13			●
<b>Smoked cigarettes on school property on at least 1 day</b> (during the 30 days before the survey)	5.3 (4.0–6.9) 1,619	5.1 (4.5–5.8) 15,851	0.85			●
<b>Ever smoked at least one cigarette every day for 30 days</b>	14.8 (11.4–19.0) 1,584	11.2 (10.0–12.6) 15,431	0.07			●
<b>Did not try to quit smoking cigarettes</b> (among students who currently smoked cigarettes, during the 12 months before the survey)	44.7 (37.7–51.9) 292	49.2 (45.9–52.6) 2,859	0.24			●
<b>Usually obtained their own cigarettes by buying them in a store or gas station</b> (among the students who were aged <18 years and who currently smoked cigarettes, during the 30 days before the survey)	18.6 (12.5–26.7) 273	14.1 (11.7–17.0) 2,266	0.23			●
<b>Used chewing tobacco, snuff, or dip on at least 1 day</b>	12.4 (9.9–15.6)	8.9 (7.3–10.8)	0.03	●		

(during the 30 days before the survey)	1,640	16,134				
<b>Used chewing tobacco, snuff, or dip on school property on at least 1 day</b> (during the 30 days before the survey)	7.8 (6.1–9.8) 1,646	5.5 (4.4–6.8) 15,974	0.03	●		
<b>Smoked cigars, cigarillos, or little cigars on at least 1 day</b> (during the 30 days before the survey)	16.7 (13.7–20.2) 1,672	14.0 (12.8–15.4) 15,671	0.12			●
<b>Smoked cigarettes; smoked cigars, cigarillos, or little cigars; or used chewing tobacco, snuff, or dip on at least 1 day</b> (during the 30 days before the survey)	26.5 (22.7–30.7) 1,519	26.0 (23.8–28.3) 14,997	0.82			●
<b>Alcohol and Other Drug Use</b>						
<b>Ever had at least one drink of alcohol on at least 1 day</b> (during their life)	70.4 (66.5–74.1) 1,588	72.5 (70.6–74.3) 15,953	0.32			●
<b>Drank alcohol for the first time before age 13 years</b> (other than a few sips)	28.9 (26.0–32.1) 1,662	21.1 (19.6–22.6) 16,207	0.00	●		
<b>Had at least one drink of alcohol on at least 1 day</b> (during the 30 days before the survey)	39.7 (35.9–43.7) 1,558	41.8 (40.2–43.4) 14,864	0.32			●
<b>Had five or more drinks of alcohol in a row within a couple of hours on at least 1 day</b> (during the 30 days before the survey)	24.6 (20.8–28.9) 1,624	24.2 (22.6–25.9) 16,009	0.87			●
<b>Usually obtained the alcohol they drank by someone giving it to them</b> (among students who currently drank alcohol during the 30 days before the survey)	—	42.2 (40.3–44.1) 6,121	~			
<b>Had at least one drink of alcohol on school property on at least 1 day</b> (during the 30 days before the survey)	6.1 (4.5–8.2) 1,625	4.5 (3.9–5.1) 16,024	0.09			●
<b>Ever used marijuana one or more times</b> (during their life)	37.4 (32.8–42.3) 1,591	36.8 (34.8–38.8) 16,060	0.80			●
<b>Tried marijuana for the first time before age 13 years</b>	9.6 (8.0–11.4) 1,602	7.5 (6.7–8.3) 16,134	0.02	●		
<b>Used marijuana one or more times</b> (during the 30 days before the survey)	17.8 (15.3–20.5) 1,600	20.8 (19.4–22.3) 16,112	0.04		●	
<b>Used marijuana on school property one or more times</b> (during the 30 days before the survey)	4.5 (2.8–7.2) 1,618	4.6 (4.0–5.4) 16,086	0.90			●
<b>Ever used any form of cocaine one or more times</b> (for example, powder, crack, or freebase, during their life)	7.5 (5.6–10.0) 1,614	6.4 (5.7–7.1) 16,204	0.32			●
<b>Used any form of cocaine one or more times</b> (for example, powder, crack, or freebase, during the 30 days before the survey)	3.8 (2.6–5.5) 1,629	2.8 (2.4–3.2) 15,557	0.18			●
<b>Ever sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times</b> (during their life)	16.8 (14.4–19.4) 1,650	11.7 (10.6–12.8) 16,293	0.00	●		
<b>Ever used heroin one or more times</b> (also called "smack", "junk", or "China white", during their life)	5.4 (3.7–8.0) 1,638	2.5 (2.2–2.9) 15,731	0.01	●		
<b>Ever used methamphetamines one or more times</b>	6.0 (4.6–7.9)	4.1 (3.6–4.6)	0.02	●		

(also called "speed", "crystal", "crank", or "ice", during their life)	1,649	16,289				
<b>Ever used ecstasy one or more times</b> (also called "MDMA", during their life)	11.0 (8.9–13.5) 1,646	6.7 (5.8–7.6) 15,887	0.00	●		
<b>Ever took steroid pills or shots without a doctor's prescription one or more times</b> (during their life)	6.4 (4.6–8.9) 1,653	3.3 (2.9–3.8) 16,320	0.00	●		
<b>Ever used a needle to inject any illegal drug into their body one or more times</b> (during their life)	4.9 (3.3–7.3) 1,638	2.1 (1.8–2.5) 16,140	0.00	●		
<b>Offered, sold, or given an illegal drug by someone on school property</b> (during the 12 months before the survey)	31.4 (28.2–34.7) 1,625	22.7 (20.7–24.9) 16,261	0.00	●		
<b>Sexual Behaviors</b>						
<b>Ever had sexual intercourse</b>	53.6 (49.4–57.7) 1,460	46.0 (42.9–49.2) 15,065	0.00	●		
<b>Had sexual intercourse for the first time before age 13 years</b>	10.3 (8.0–13.1) 1,471	5.9 (5.1–6.8) 15,046	0.00	●		
<b>Had sexual intercourse with four or more persons</b> (during their life)	18.0 (14.8–21.7) 1,464	13.8 (12.4–15.4) 14,987	0.03	●		
<b>Had sexual intercourse with at least one person</b> (during the 3 months before the survey)	38.9 (34.6–43.3) 1,460	34.2 (31.9–36.5) 15,017	0.06			●
<b>Drank alcohol or used drugs before last sexual intercourse</b> (among students who were currently sexually active)	20.1 (15.6–25.6) 494	21.6 (20.0–23.3) 5,548	0.57			●
<b>Did not use a condom during last sexual intercourse</b> (among students who were currently sexually active)	41.3 (34.5–48.5) 476	38.9 (36.9–41.0) 5,454	0.50			●
<b>Did not use birth control pills before last sexual intercourse</b> (to prevent pregnancy, among students who were currently sexually active)	80.6 (74.8–85.3) 478	80.2 (77.5–82.6) 5,360	0.89			●
<b>Were never taught in school about AIDS or HIV infection</b>	17.6 (15.0–20.6) 1,580	13.0 (11.7–14.3) 16,109	0.00	●		
<b>Did not use Depo-Provera before last sexual intercourse</b> (to prevent pregnancy, among students who were currently sexually active)	95.1 (92.6–96.7) 478	96.9 (96.3–97.4) 5,360	0.07			●
<b>Did not use birth control pills or Depo-Provera before last sexual intercourse</b> (to prevent pregnancy, among students who were currently sexually active)	75.6 (69.6–80.8) 478	77.1 (74.3–79.7) 5,360	0.63			●
<b>Did not use both a condom during last sexual intercourse and birth control pills or Depo-Provera before last sexual intercourse</b> (to prevent pregnancy, among students who were currently sexually active)	89.9 (85.9–92.8) 469	91.1 (89.7–92.4) 5,329	0.49			●
<b>Dietary Behaviors</b>						
<b>Ate fruits and vegetables less than five times per day</b> (100% fruit juices, fruit, green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	85.1 (82.3–87.5) 1,566	77.7 (76.3–78.9) 16,057	0.00	●		

<b>Did not drink 100% fruit juices</b> (during the 7 days before the survey)	26.1 (23.0–29.5) 1,600	19.4 (18.0–20.8) 16,167	0.00	●		
<b>Did not eat fruit</b> (during the 7 days before the survey)	17.9 (15.4–20.7) 1,620	11.4 (10.1–12.9) 16,232	0.00	●		
<b>Did not eat green salad</b> (during the 7 days before the survey)	40.4 (36.3–44.6) 1,620	36.6 (34.7–38.6) 16,220	0.09			●
<b>Did not eat potatoes</b> (excluding French fries, fried potatoes, or potato chips, during the 7 days before the survey)	32.9 (29.8–36.1) 1,613	31.2 (29.5–33.1) 16,224	0.36			●
<b>Did not eat carrots</b> (during the 7 days before the survey)	60.7 (57.9–63.4) 1,612	51.8 (49.5–54.2) 16,223	0.00	●		
<b>Did not eat other vegetables</b> (excluding green salad, potatoes, or carrots, during the 7 days before the survey)	23.8 (20.7–27.2) 1,613	16.7 (15.3–18.1) 16,215	0.00	●		
<b>Drank a can, bottle, or glass of soda or pop at least one time per day</b> (not including diet soda or diet pop, during the 7 days before the survey)	33.5 (29.2–38.1) 1,602	29.2 (27.2–31.2) 16,188	0.07			●
<b>Drank less than three glasses per day of milk</b> (during the 7 days before the survey)	91.5 (89.6–93.1) 1,608	85.5 (83.1–87.6) 16,174	0.00	●		
<b>Ate fruit or drank 100% fruit juices less than two times per day</b> (during the 7 days before the survey)	74.3 (70.5–77.7) 1,593	66.1 (64.4–67.8) 16,141	0.00	●		
<b>Ate vegetables less than three times per day</b> (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	88.3 (85.8–90.5) 1,595	86.2 (85.2–87.1) 16,160	0.09			●
<b>Overweight</b> (students who were $\geq$ 85th percentile but $<$ 95th percentile for body mass index, by age and sex, based on reference data)	15.7 (13.3–18.6) 1,483	15.8 (14.7–17.0) 15,178	0.96			●
<b>Obese (students who were <math>\geq</math> 95th percentile for body mass index, by age and sex, based on reference data)</b>	14.4 (12.1–17.2) 1,483	12.0 (10.9–13.1) 15,178	0.07			●
<b>Described themselves as slightly or very overweight</b>	27.3 (24.0–30.9) 1,626	27.7 (26.7–28.6) 16,123	0.85			●
<b>Did not exercise to lose weight or to keep from gaining weight</b> (during the 30 days before the survey)	41.4 (38.7–44.1) 1,603	38.5 (37.2–39.8) 16,109	0.05			●
<b>Did not eat less food, fewer calories, or low-fat foods to lose weight or to keep from gaining weight</b> (during the 30 days before the survey)	59.8 (56.7–62.8) 1,599	60.5 (59.0–61.8) 16,099	0.68			●
<b>Went without eating for 24 hours or more to lose weight or to keep from gaining weight</b> (during the 30 days before the survey)	16.6 (14.2–19.3) 1,600	10.6 (9.7–11.5) 16,115	0.00	●		
<b>Took diet pills, powders or liquids to lose weight or to keep from gaining weight</b> (without a doctor's advice, during the 30 days before the survey)	10.5 (8.4–13.2) 1,601	5.0 (4.5–5.5) 16,120	0.00	●		
<b>Vomited or took laxatives to lose weight or to keep from</b>	9.3 (6.8–12.5)	4.0 (3.5–4.4)	0.00	●		

<b>gaining weight</b> (during the 30 days before the survey)	1,590	16,100				
<b>Physical Activity</b>						
<b>Physically active at least 60 minutes per day on less than 5 days</b> (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	58.0 (53.6–62.4) 1,610	63.0 (61.2–64.8) 16,109	0.03		●	
<b>Did not attend physical education classes in an average week</b> (when they were in school)	60.6 (54.9–66.0) 1,579	43.6 (36.4–51.1) 15,578	0.00	●		
<b>Did not attend physical education classes daily</b> (when they were in school)	77.3 (72.5–81.5) 1,579	66.7 (60.7–72.3) 15,578	0.00	●		
<b>Did not play on sports teams</b> (run by their school or community groups during the 12 months before the survey)	47.8 (44.0–51.6) 1,553	41.7 (38.9–44.5) 15,562	0.01	●		
<b>Watched television 3 or more hours per day</b> (on an average school day)	36.4 (31.9–41.0) 1,588	32.8 (30.4–35.3) 16,126	0.17			●
<b>Used computers 3 or more hours per day</b> (played video or computer games or used a computer for something that was not school work on an average school day)	21.0 (18.2–24.2) 1,578	24.9 (22.9–27.0) 16,124	0.03		●	
<b>Physically active at least 60 minutes per day on less than 7 days</b> (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	75.7 (73.2–78.0) 1,610	81.6 (80.5–82.7) 16,109	0.00		●	
<b>Did not participate in at least 60 minutes of physical activity on any day</b> (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	19.5 (16.7–22.6) 1,610	23.1 (21.5–24.8) 16,109	0.03		●	
<b>Asthma</b>						
<b>Ever told by a doctor or nurse that they had asthma</b>	23.2 (21.0–25.6) 1,584	22.0 (20.8–23.1) 16,089	0.33			●
<b>Ever told by a doctor or nurse that they had asthma and still have asthma</b>	9.9 (8.0–12.2) 1,572	10.8 (9.9–11.7) 15,998	0.42			●

**Footnotes**

†	Percentage, confidence interval, cell size
‘—’	= Data not available
~	= P-value not available

**Application URL:** <http://ngdev.cdc.gov/YouthOnline/App/Results.aspx?TT=G&OUT=0&SID=HS&QID=QQ&LID=AR&YID=2009&LID2=XX&YID2=2009&COL=&ROW1=&ROW2=&HT=QQ&LCT=&FS=1&FR=1&FG=1&FSL=&FRL=&FGL=&PV=&TST=True&C1=AR2009&C2=XX2009&QP=G&DP=1&VA=CI&CS=Y&SYID=&EYID=&SC=DEFAU LT&SO=ASC&PF=1>

